



Animal Nutrition
 1735 Market Street
 Philadelphia, PA 19103
 800 356 4657
Wendy.plocher@tronox.com

Feeding Recommendations for S-Carb® in Poultry

Broilers:

Expected benefits from S-Carb addition are improved tolerance to heat stress, improved performance, and drier litter.

Basal Addition: Replace 5 lbs/ton of Salt with 5 lbs/ton of S-Carb

This replacement will: Lower Cl 0.15% (42 meq/kg)
 Lower Na 0.02% (9 meq/kg)
 & raise DEB 33 meq/kg

During Heat Stress: Replace up to 8 lbs of Salt with S-Carb.

This replacement will: Lower Cl 0.24% (68 meq/kg)
 Lower Na 0.03% (13 meq/kg)
 & raise DEB 55 meq/kg

***** Recommended DEB is 200-300 meq/kg *****

Turkeys:

Use S-Carb to replace salt when dietary chloride is high due to inclusion of ingredients with high chloride contents, such as lysine hydrochloride or choline chloride. S-Carb will help maintain sodium without added chloride.

Layers:

For Older birds or all layers during heat stress to improve eggshell quality.

Replace 6 lbs/ton of Salt with 6 lbs/ton of S-Carb

This replacement will: Lower Cl 0.18% (51 meq/kg)
 Lower Na 0.02% (9 meq/kg)
 & raise DEB 42 meq/kg

Always insure that minimum dietary chloride is met.

Minimum Chloride Requirements:

	Starter	Grower	Finish
Broiler	0.20 %	0.15 %	0.12 %
Turkey	0.17 %	0.15 %	0.12 %

Layer	0.11-0.16%
-------	------------

NRC 1994